

Relative coherent modules and semihereditary modules

Lixin Mao

Department of Mathematics and Physics, Nanjing Institute of Technology

Email: maolx2@hotmail.com

Given a positive integer n , a left R -module M is called n -coherent (resp. n -semihereditary) if every n -generated submodule of M is finitely presented (resp. projective). We investigate the properties of n -coherent modules and n -semihereditary modules. Various results are developed, many extending known results.

REFERENCES

1. J.L. Chen, N.Q. Ding, *A note on existence of envelopes and covers*, Bull. Austral. Math. Soc. **54** (1996), 383-390.
2. J.L. Chen, N.Q. Ding, Y.L. Li, Y.Q. Zhou, *On (m, n) -injectivity of modules*, Comm. Algebra **29** (2001), 5589-5603.
3. G. Dai, N. Ding, *Coherent rings and absolutely pure covers*, Comm. Algebra **46** (2018), 1267-1271.
4. E.E. Enochs, O.M.G. Jenda, *Relative Homological Algebra*, Walter de Gruyter, Berlin-New York, 2000.
5. A. Shamsuddin, *n -injective and n -flat modules*, Comm. Algebra, **29** (2001), 2039-2050.
6. X.X. Zhang, J.L. Chen, *On n -semihereditary and n -coherent rings*, Inter. Electronic J. Algebra **1** (2007), 1-10.
7. X.X. Zhang, J.L. Chen, J. Zhang, *On (m, n) -injective modules and (m, n) -coherent rings*, Algebra Colloq. **12** (2005), 149-160.
8. Z.M. Zhu, Z.S. Tan, *On n -semihereditary rings*, Sci. Math. Jpn. **62** (2005), 455-459.